

Appendix D

PowerPoint Presentation
prepared by Tetra Tech:

“Winter Manure Application and Water
Quality; Overview of the Literature”

Oct. 30, 2014



Winter Manure Application and Water Quality

Overview of the Literature



Overview of Briefing

- Define winter application and applicability
- Why winter application can occur
- Research findings on winter application water quality impacts and risks
- Management options
- Research needs
- Conclusions

Purpose of White Paper

Survey of existing research on winter manure application and water quality to support an informed discussion on winter application and potential options for mitigating the effects of unavoidable winter application.



Process

All sources of research data, management recommendations, and other information concerning winter application of animal manure:

- SERA-17 bibliography <http://www.sera17.ext.vt.edu/index.htm>.
- On-line databases: NAL, Google Scholar
- JEQ, JSWC, Trans. ASABE, others
- Grey literature
- Extension publications



Complete list of references in appendix of this presentation




Definitions & Applicability

Regions that experience frozen and/or snow-covered soils face special constraints for good manure management.

Historical snowfall and soil frost data suggest that winter application practices warrant careful consideration in parts or all of U.S. states except Florida, Hawaii, Louisiana, and Mississippi.

Annual Mean Total Snowfall



Definitions & Applicability

Land application of animal manure to snow-covered, ice-covered, or frozen soils.

Other situations outside the scope of the paper where risks also exist:

- “Warm winter” application during high precipitation periods
- Application to saturated soils at any time
- Others....



Agronomic benefits of winter application?

The comprehensive literature review found no published research to support agronomic factors as a basis for recommending winter manure application:

- Prevention of soil compaction*
- Reduction of N volatilization*
- Providing crop nutrients for spring planting,
- Reduce risk of excessive manure application around a ban period

* Both have been documented as “positives” of winter application, but other factors outweigh the benefits



Winter Application Impacts and Risks

Hydrologic factors

- **Frozen soils decrease infiltration and increase runoff**
 - Most (not all) frozen soils virtually impervious (Fleming and Fraser 2000)
 - 56% increase in runoff volume from frozen soils (Williams et al. 2012)
 - Catchments with frozen soils have greater water yield than unfrozen soils (Molnau and Cherry 1990)



Winter Application Impacts and Risks

Hydrologic factors

- **Importance of snowmelt in annual discharge**
 - WI: 50% of annual ag runoff in snowmelt (Stuntebeck et al. 2011)
 - Alberta: 90% of annual runoff (Little et al. 2007)
 - Sask.: 85% of annual runoff and 50% of groundwater recharge (Maule and Elliott (2005a)



Winter Application Impacts and Risks

Hydrologic factors

- **Critical characteristics**
 1. Structure of the frozen soil
 2. Depth of penetration of the frost
 3. Persistence of soil frost
 4. Areal extent of frozen soil. (Storey 1955)
- Freeze-thaw processes poorly understood (Storey 1955)
- Function of variable soil and climate characteristics, e.g., tillage, cover, moisture (Storey 1955, Willis et al. 1961)



Winter Application Impacts and Risks

- Frozen soils and snowpack increase the risk of runoff from winter-applied manure.
- Dormant or absent crops provide no nutrient uptake
- Incorporation difficult or impossible
- Freezing does not reliably kill pathogens
- Loss of soluble nutrients and microorganisms



Winter Application Impacts and Risks

Research data demonstrate that substantial potential exists for runoff losses of N and P and impacts to receiving waters if manure is applied to frozen soils or snow-covered ground.



- Very high nutrient concentrations, e.g.,
 - TP 1.6 – 15.4 mg/L; TKN 24 – 1086 mg/L
 - TP increase 165 – 224%; soluble P increase 246 – 1480%; 576% increase in NH₃-N in runoff following winter application of dairy manure
- Mass losses of up to 27% of applied P, 22% of applied N

Thompson et al. 1979, Melvin and Lorimer 1996, Phillips et al. 1981, Clausen 1990, 1991, Midgeley and Dunklee 1945, Hensler et al. 1970, Phillips et al. 1975, Converse et al. 1976, Klausner et al. 1976, Young and Mutchler 1976



Winter Application Impacts and Risks

The magnitude of nutrient losses from winter-applied manure appears to be controlled by a large number of factors whose relative influence is poorly understood, including:

- Effects of soil frost on infiltration and runoff generation;
- Timing, extent, and depth of snow cover relative to manure application;
- Soil temperature;
- Snowmelt and winter rainfall;
- Timing of manure application relative to snowfall, snowmelt, and runoff;
- The form of manure applied;
- Land use/land cover, including crop, crop residue, and tillage;
- Land slope;
- Existing conservation practices; and
- Weather and climate.



Winter Application Impacts and Risks


Most of the major factors that appear to influence nutrient losses from winter-applied manure are highly variable and exceptionally difficult to predict with any certainty on a fine scale.

→ Even if all the processes governing nutrient losses from winter-applied manure are perfectly understood, fully avoiding adverse water quality impacts – or even reasonable estimation of the risk of significant water quality impacts – from winter manure application on a site-specific basis will continue to be extremely challenging.



Winter Application Impacts and Risks

Although the potential for major nutrient losses is not always realized, there is ample evidence in the literature that runoff losses of N and P from winter-applied manure are often significant, both in terms of agronomic losses and potential water quality impacts.



 TETRATECH

Management standards

Some states have adopted technical standards for CAFOs, but these are highly variable

Documented in Appendix to White Paper

State	Type of Restriction	Restriction Details		Mechanism and Source
		Based on Form-Defined Emergency Conditions Only	Based on Emergency Conditions and Incidents	
Pennsylvania	✓	<p>Applicable to NYSDEC-permitted CAFOs.</p> <p>Manure may not be spread during the winter (December 15 through February 28); or any time the ground is frozen at least four (4) inches, or any time that the ground is covered by snow or ice.</p> <p>Permittees must have a Winter Period Management Plan, or unless the permittee has fully complied with 27 Pa. Code § 133.33(d) and (e) and (f) for all applicable circumstances. The permittee shall notify DEP in writing within 7 days prior to the start of the winter period.</p> <p>The permittee shall certify in the notification that:</p> <ol style="list-style-type: none"> Consistent with the approved NMP, actions have been taken to minimize the risk of spreading manure or sludge onto storage facilities prior to the winter period to prevent pollution of ground and surface water; The manure or sludge is applied in accordance with the NMP; Manure spread during the winter shall be applied in an amount and a manner that minimizes the risk of surface water pollution. <p>The permittee may use the Department of Environmental Protection, Winter Period Application of Manure Notification form to provide the notification.</p> <p>There shall be no winter application:</p> <ol style="list-style-type: none"> Within 100 feet of an above-ground agricultural drain inlet where surface water runoff may collect; Within 100 feet of a wetland on the National Wetland Inventory maps which is within the 100-year floodplain of an EV reaches segment if surface water runoff may collect; Unless the fields have 25% cover or an established cover crop... 	<p>Applicable to all operations.</p> <p>590 standard</p>	

 TETRATECH

BMPs

Most cold-climate states and provinces recommend against winter manure application and consider winter spreading to be a last resort if storage is limited or if weather impairs recommended management



BMPs

Some management measures may be required to mitigate emergency or unavoidable winter application

There is currently no body of standards and specifications supported by research data for BMPs or other management measures to specifically mitigate potential impacts of winter manure application.



BMPs

Vegetation-based practices are largely dormant and less effective during critical mid-winter thaw and spring runoff periods when most nutrient loss occurs



BMPs

Common sense recommendations like increased setbacks or reduced application rates may have some effect on reducing runoff losses of nutrients from winter-spread manure, but there is little documentation in the literature.

- Setbacks
- Avoid flood zones
- Slope criteria
- Proximity to water
- Residue cover
- Reduced application rate
- Incorporation/injection



Risk-based approaches

Check Wisconsin's Online Runoff Risk Advisory Forecast
www.manureadvisorysystem.wi.gov



If it's RED Don't SPREAD
 Working together to keep manure out of lakes and streams

Runoff Risk Advisory Forecast
 Wisconsin Manure Management Advisory System

[MMAS Home](#) | [500 Nutrient Appl. Restriction Maps](#) | [Runoff Risk Advisory Forecast Map](#) | [InteractiveOn](#)

Runoff Risk Advisory Forecast Snapshot

This forecast was valid from approximately Mar 08 2014 09:00AM until it was updated with a more recent model run at Mar 08 2014 07:30PM

Runoff Risk (3-day)	Winter Risk (10 day)
Low	Moderate
Moderate	High
High	

Legend:

- Frozen/snow-covered soil
- Runoff
- Snowmelt Runoff



Important assumptions must be communicated to users:

- This is *not* meant to be the *only* tool for deciding when to spread
- This approach will never produce perfect prediction
 - One farm may have runoff, the next one may not
 - Rainfall patterns, snowpack conditions, field aspect, etc.
- Users must *combine* knowledge of local conditions with forecast

<http://www.manureadvisorysystem.wi.gov/>

TETRATECH

Risk-based approaches

New York:

- Identify the lowest risk fields for spreading as a last resort (e.g., when storage is full)
- Evaluate runoff potential along with other management needs: soil wetness, weather forecast for rainfall or snowmelt, presence of diversions or field ditches and drainage tile, rate per acre, and total amount of manure to be applied.
- When conditions for runoff are high, consider delaying the application, reducing the rate, reducing the total amount applied, and/or applying smaller amounts of manure over a period of days rather than hours
- Avoid application when:
 - Significant rainfall or snowmelt is predicted within 24-48 hours.
 - Soil is frozen, snow covered or saturated
 - Tile drains are flowing from field drainage

Czymmek et al. (2005)

TETRATECH


Risk-based approaches

Ontario:

Risk assessment of conditions unsuitable for manure application:

- Soil frost depth >0.05 m
- Snow accumulation >0.05 m
- Soil volumetric water content \geq plastic limit

Range of recommended dates for spring start and fall end of manure application based on probabilities of one or more of the limiting criteria



Fallow et al. (2007)



Many unknowns about BMPs

- ▶ Research conflicting on effectiveness of slope restrictions or requirements for vegetative cover;
- ▶ Insufficient experience with winter manure incorporation or injection to recommend a BMP
- ▶ Runoff control measures may be effective, but trade-offs with leaching are unknown;
- ▶ Effectiveness of VFS or buffers in capturing or treating runoff from winter application is uncertain because snowmelt/runoff occurs when vegetative measures are essentially dormant.
- ▶ Net effect of risk-based procedures un-tested



BMP Research needs

- ▶ Identify, quantify, and prioritize individual factors influencing nutrient and pathogen losses from winter-applied manure
- ▶ Assess the real risks of major nutrient losses when limited storage and winter manure application restrictions promote high manure applications during the time immediately before or after a ban period.
- ▶ Document the effectiveness of vegetative BMPs on delivery of nutrients from winter-manured fields to surface waters during the mid-winter thaw and spring snowmelt periods.
- ▶ Evaluate the effects of BMPs currently recommended by NRCS for year-round implementation to reduce surface runoff losses of nutrients and pathogens from winter manure application and on leaching losses, in particular through subsurface drainage.
- ▶ Field test and evaluate the practicality and effectiveness of agricultural implements designed to incorporate manure directly into frozen and/or snow-covered soils.



 TETRATECH

Weight of Evidence

- Lack of agronomic benefit,
- Documented water quality impacts, and
- Absence of effective BMPs



Avoid winter
manure
application



 TETRATECH



QUESTIONS?



References

Balkcom, K.S., A.M. Blackmer, D.J. Hansen, T.F. Morris, and A.P. Mallarino. 2003. Testing Soils and Cornstalks to Evaluate Nitrogen Management on the Watershed Scale. *J. Environ. Qual.* 32:1015–1024.

Baker, D.G. 1972. Prediction of spring runoff. *Water Resour. Res.* 8:966-972.

Brenneman, G. undated. Manure Manager Series, Vol. 3. Iowa State University Extension. <http://www.agronext.iastate.edu/immag/pubs/imms/vol3.pdf> (Accessed 9-25-2013).

Bishop, P.L., W.D. Hively, J.R. Stedinger, M.R. Rafferty, J.L. Lojperberger, and J.A. Bloomfield. 2005. Multivariate analysis of paired watershed data to evaluate agricultural best management practice effects on stream water phosphorus. *J. Environ. Qual.* 34:1087–1101.

Brown, M.B., P. Longabucco, M.R. Rafferty, P.D. Robillard, M.F. Walter, and D.A. Haith. 1989. Effects of animal waste control practices on nonpoint source phosphorus loading in the West Branch of the Delaware River watershed. *J. Soil Water Cons.* 44(1):67-70.



References

Carrington, E.G. and M.E. Ransome. 1994. Factors influencing the survival of *Cryptosporidium* oocysts in the environment. Report No. FR 0456. *Foundations for Water Research*. Marlow, Bucks.

Clausen, J.C. 1990. Winter and Fall application of manure to corn land. Pages 179 – 180 in Meals, D.W. 1990. LaPlatte River Watershed Water Quality Monitoring and Analysis Program: Comprehensive Final Report. Program Report No. 12. Vermont Water Resource Research Center, University of Vermont, Burlington.

Clausen, J.C. 1991. Best manure management effectiveness. Pages 193 – 197 in Vermont RCWP Coordinating Committee. 1991. St. Albans Bay Rural Clean Water Program, Final Report. Vermont Water Resources Research Center, University of Vermont, Burlington

Clausen, J.C. and D.W. Meals. 1989. Water quality achievable with agricultural best management practices. *J. Soil Water Cons.* 44(6):593-596.

Converse, J.C., G.D. Bubenzer, and W.H. Paulson. 1976. Nutrient losses in surface runoff from winter spread manure. *Trans. ASAE* 19:517-519.



References

Cooley, K.R. (ed.), 1990. Proceedings of an International Symposium Frozen Soil Impacts on Agricultural, Range, and Forest Lands. Special Report 90-1, U.S. Army Cold Regions Research and Engineering Laboratory, Hanover, NH.

Dunne, T. 1983. Relation of field studies and modeling in the prediction of storm runoff. *J. Hydrol.* 65:25-48.

Fallow, D.J., D.M. Brown, J.D. Lauzon, and G.W. Parkin. 2007. Risk assessment of unsuitable winter conditions for manure and nutrient application across Ontario. *J. Environ. Qual.* 36:31–43

Fayer, R. and T. Nerad. 1996. Effects of low temperature on viability of *Cryptosporidium parvum* oocysts. *Appl. and Environ. Microbiol.* 62(4):1431-1433

Fleming, R. and H. Fraser. 2000. Impacts of Winter Spreading of Manure on Water Quality - Literature Review. University of Guelph, Report prepared for Ontario Pork, Etobicoke, ON, Canada.



References

Gerba, D., C. Wallis, and J. Mellnick. 1975. Fate of wastewater bacteria and viruses in soil. *J. Irr. Drain. Div. ASCE* 101:157-174.

Gessel, P. D., N. C. Hansen, J. F. Moncrief, and M. A. Schmitt. 2004. Rate of fall-applied liquid swine manure: effects on runoff transport of sediment and phosphorus. *J. Environ. Qual.* 33:1839-1844.

Gilley, J.E., L.M. Risso, and B. Eghball. 2002. Managing runoff after manure application. *J. Soil Water Cons.* 57(6):530-533.

Hansen, N.C., S.C. Gupta, and J.F. Moncrief. 2000. Snowmelt runoff, sediment, and phosphorus losses under three different tillage systems. *Soil Tillage Res.* 57:93-100.

Hensler, R.F., R.J. Olsen, S.A. Witzel, O.J. Attoe, W.H. Paulson, and R.F. Johannes. 1970. Effect of method of manure handling on crop yields, nutrient recovery and runoff losses. *Trans. ASAE*:726-731.



References

Hernandez, J.A. and M.A. Schmitt. 2012. Manure Management in Minnesota. University of Minnesota Extension. <http://www.extension.umn.edu/distribution/cropsystems/DC3553.html> (Accessed 9-25-2013).

Jann, S.M. 2005. Interim Final Technical Guidance for the Application of CAFO Manure on Land in the Winter. United States Environmental Protection Agency, Region 5, Chicago, IL. http://www.cals.ncsu.edu/waste_mgt/natcenter/sanantonio/Jann.pdf (Accessed 9-24-2013).

Kibby, H.J., C. Hagedorn, and E.L. McCoy. 1978. Use of Fecal Streptococci as indicators of pollution in soil. *Appl. and Environ. Microbiol.* 35(4):711-717.

Klausner, S.D., Zwerman, P.J., and Ellis, D.F. 1976. Nitrogen and phosphorus losses from winter disposal of manure. *J. Environ. Qual.* 5(1):47-49



References

Klausner, S.D., Zwerman, P.J., and Ellis, D.F. 1976. Nitrogen and phosphorus losses from winter disposal of manure. *J. Environ. Qual.* 5(1):47-49

Komiskey, M.J., T.D. Stuntebeck, D.R. Frame, and F.W. Madison. 2011. Nutrients and sediment in frozen-ground runoff from no-till fields receiving liquid-dairy and solid-beef manures. *J. Soil Water Cons.* 66(5):303-312.

Kongoli, C.E. and W.L. Bland. 2002. Influence of manure application on surface energy and snow cover. *J. Environ. Qual.* 31:1166-1173.

Kongoli, C.E. and W.L. Bland. 2002a. Influence of manure application on surface energy and snow cover: Field experiments. *J. Environ. Qual.* 31:1166-1173.

Kudva, I.T., K. Blanch, and C.J. Hovde. 1998. Analysis of *Escherichia coli* O157:H7 in ovine or bovine manure and manure slurry. *Appl. and Environ. Microbiol.* 64(9):3166-3174.



References

Lauer, D.A., D.R. Bouldin, and S.D. Klausner. 1976. Ammonia volatilization from dairy manure spread on the soil surface. *J. Environ. Qual.* 5:134-141.

Lemunyon, J.L., and R.G. Gilbert. 1993. The concept and need for a phosphorus assessment tool. *J. Prod. Agric.* 6:483-486.

Lewis, T.W. and J.C. Makarewicz. 2009. Winter application of manure on an agricultural watershed and its impact on downstream nutrient fluxes. *J. Great Lakes Res.* 35(sp1):43-49.

Little, J.L., S.C. Nolan, J.P. Casson, and B.M. Olson. 2007. Relationships between Soil and Runoff Phosphorus in Small Alberta Watersheds. *J. Environ. Qual.* 36:1289-1300.

Lorimer, J.C. 1999. Minimizing risk when applying manure in winter. Iowa State University Extension, Ames, IA.
<http://www.extension.iastate.edu/Pages/communications/EPC/F99/winter.html>

Lorimer, J.C. and J.C. Melvin. 1996. Nitrogen losses in surface runoff from winter-applied manure. University of Iowa. Final Report.



References

Madison, F., K. Kelling, L. Massie, and L.W. Good. 2003. *Guidelines for applying manure to cropland and pasture in Wisconsin*. University of Wisconsin Extension. <http://learningstore.uwex.edu/assets/pdfs/A3392.pdf> (Accessed 9-25-2013).

Marsh, P. 1999. Snowcover formation and melt: recent advances and future prospects. *Hydrol. Proc.* 13:2117-2134.

Maulé, C. and J. Elliott. 2005a. Effect of Hog Manure Injection Upon Soil Productivity and Water Quality; Part I Perdue Site, 1998-2004. ADF Project 98000094. Dept of Agricultural and Bioresource Engineering, University of Saskatchewan, 57 Campus Drive, Saskatoon, SK.

McCool, D.K. 1990. Crop management effects on runoff and soil loss from thawing soil. Pp. 171-176 *In: Frozen soil impacts on agricultural, range, and forest lands*. K.R. Cooley (ed.) Proceedings of International Symposium, Spokane, Washington. CRREL Special Report 90-1. Cold Regions Research and Engineering Laboratory. U.S. Army Corps of Engineers, Hanover, NH.



References

Melvin, S. and J. Lorimor. 1996. Effects of winter manure spreading on surface water quality. 1996 Research Report, Agricultural & Biosystems Engineering, Iowa State University Extension, Ames, IA. <http://www.nppc.org/Research/96Reports/96Melvin-manure.html>

Michigan Commission of Agriculture & Rural Development. 2013. Generally Accepted Agricultural and Management Practives for Manure Management and Utilization. Lansing, MI. http://www.michigan.gov/documents/mdard/2013_FINAL_MANURE_GAAMP_409764_7.pdf

Midgeley, A.R. and D.E. Dunklee. 1945. Fertility runoff losses from manure spread during the winter. Univ. of Vermont, Agric. Exp. Station, Bulletin 523, 19 p.

Molnau, M. and J.G. Cherry. 1990. A comparison of runoff occurring on frozen and unfrozen soils. Pp. 279-281 *In: Frozen soil impacts on agricultural, range, and forest lands*. K.R. Cooley (ed.) Proceedings of International Symposium, Spokane, Washington. CRREL Special Report 90-1. Cold Regions Research and Engineering Laboratory. U.S. Army Corps of Engineers, Hanover, NH.



References

Moore, I.C. and F.W. Madison. 1985. Description and application of an animal waste phosphorous loading model. *J. Environ. Qual.* 14(3):364-368.

Nolan, S., L. Good, P. Loro, J. Elliot, T. Wallace, and B. Olson. Undated. Best Management Practices for Snowmelt Runoff. Alberta Agriculture and Rural Development. Edmonton, AB.

Ohio State University. 2011. *Properly Applying Manure on Frozen Ground*. OSU Crop Observation and Recommendation Network (C.O.R.N.) Newsletter 2011-42 <http://corn.osu.edu/newsletters/2011/2011-42/properly-applying-manure-on-frozen-ground> (Accessed 9-25-2013).

Ontario Ministry of Agriculture and Food. 2011. Winter Application of Manure and Other Agricultural Source Materials. OMAFRA Fact Sheet 10-073. <http://www.omafra.gov.on.ca/english/engineer/facts/10-073.htm#5> (Accessed 9-25-2013).

Owens, L.B., J.V. Bonta, M.J. Sipitalo, and S. Rogers. 2011. Effects of winter manure application in Ohio on the quality of surface runoff. *J. Environ. Qual.* 40:153-165.



References

Pappas, E.A., R.S. Kanwar, J.L. Baker, J.C. Lorimor, and S. Mickelson. 2008. Fecal indicator bacteria in subsurface drain water following swine manure application. *Trans. ASABE* 51:1567-1573.

Patni, N.K., H.R. Toxopeus, and P.Y. Jui. 1985. Bacterial quality of runoff from manured and non-manured cropland. *Trans. ASAE* 28():1871-1884.

Phillips, P.A., A.J. MacLean, F.R. Hore, F.J. Sowden, A.D. Tenant, and N.K. Patni. 1975. Soil water and crop effects of selected rates and times of dairy cattle liquid manure applications under continuous corn. Engineering Research Service Contribution No. 540. Agriculture Canada, Ottawa, Ontario.

Philips, P.A., Culley, J.L.B., Hore, F.R., and Patni, N.K. 1981. Pollution potential and corn yields from selected rates and timing of liquid manure applications. *Trans. ASAE*. 1981: 139-144

Pikul, J.L., Jr., J.F. Zuzel, and D.E. Wilkins. 1992. Infiltration into frozen soil as affected by ripping. *Trans. ASAE*. 35:83-90.



References

Pikul, J.L., Jr., D.E. Wilkins, J.K. Aase, and J.F. Zuzel. 1996. Contour ripping: A tillage strategy to improve water infiltration into frozen soil. *J. Soil Water Cons.* 51:76-83.

Pionke, H.B., W.J. Gburek, R.R. Schnabel, A.N. Sharpley, and G.F. Elwinger. 1999. Seasonal flow, nutrient concentrations and loading patterns in stream flow draining an agricultural hill-land watershed. *J. Hydrol.* 220:62-73.

Reddy, K.R., R. Khaleel, and M.R. Overcash. 1981. Behavior and transport of microbial pathogens and indicator organisms in soils treated with organic wastes. *J. Environ. Qual.* 10(3):255-266.

Ricker, D.H. and J.D. Smolik. 1990. Tillage and crop residue effects on soil frost depth. Pages 31-35 IN Proceedings of an International Symposium Frozen Soil Impacts on Agricultural, Range, and Forest Lands. Cooley, K.R. (ed.), Special Report 90-1, U.S. Army Cold Regions Research and Engineering Laboratory, Hanover, NH.



References

Schillinger, W.F., and D.E. Wilkins. 1997. Deep ripping fall-planted wheat after fallow to improve infiltration and reduce erosion. *J. Soil. Water Cons.* 52:198-202.

Sharpley, A.N. and P. Kleinman. Effect of rainfall simulator and plot scale on overland flow and phosphorus transport. *J. Environ. Qual.* 32(6):2172-2179.

Srinivasan, M.S.. R.B. Bryant, M.P. Callahan, and J.L. Weld. 2006. Manure management and nutrient loss under winter conditions: A literature review. *J. Soil Water Cons.* 61(4):200-209.

Steenhuis, T.S., G.D. Bubenz, and J.S. Converse. 1979. Ammonia volatilization of winter spread manure. *Trans. ASAE* 22: 153-157.

Steenhuis, T.S., G.D. Bubenz, J.C. Converse, and M.F. Walter. 1981. Winter-spread Manure Nitrogen loss. *Trans. ASAE* 24(2):436-441

Stoddard, C.S., M.S. Coyne, and J.H. Grove. 1998. Fecal bacteria survival and infiltration through a shallow agricultural soil: timing and tillage effects. *J. Environ. Qual.* 27:1516-1523.



References

Storey, H.C. 1955. Frozen soil and spring and winter floods. Pp. 179-185. *In: The Yearbook of Agriculture*. U.S. Government Printing Office, Washington, D.C.

Stuntebeck, T.D., M.J. Komiskey, M.C. Peppler, D.W. Owens, and D.R. Frame. 2011. Precipitation-runoff relations and water-quality characteristics at edge-of-field stations, Discovery Farms and Pioneer Farm, Wisconsin, 2003–8: U.S. Geological Survey Scientific Investigations Report 2011–5008, 46 p.

Thompson, D.B., T.L. Loudon, and J.B. Gerrish. 1979. Animal manure movement in winter runoff for different surface conditions. Pages 145-157 *In* R.C. Loehr et al., eds. Best Management Practices for Silviculture and Agriculture. Ann Arbor Science, Ann Arbor, MI.

Ulen, B. 2003. Concentrations and transport of different forms of phosphorus during snowmelt runoff from an illite clay soil. *Hydrol. Proc.* 17:747-758.

U.S. EPA. 2012. *NPDES Permit Writers' Manual for Concentrated Animal Feeding Operations*. EPA 833-F-12-001. U.S. Environmental Protection Agency, Office of Water. Washington, DC.



References

Vansteelant, J.Y. 2000. Personal communication, Institut National de la Recherche Agronomique, Thonon les Bains, France.

Williams, M.R., G.W. Feryereisen, D.B. Beegle, and R.D. Shannon. 2012a. Soil temperature regulates phosphorus loss from lysimeters following fall and winter manure application. *Trans. ASABE* 55(3):871-880.

Williams, M.R., G.W. Feryereisen, D.B. Beegle, and R.D. Shannon. 2012b. Soil temperature regulates nitrogen loss from lysimeters following fall and winter manure application. *Trans. ASABE* 55(3):861-870.

Willis, W.O., C.W. Carlson, J. Alessi, and H.H. Haas. 1961. Depth of freezing and spring runoff as related to fall soil-moisture level. *Can. J. Soil Sci.* 41:115-123.

Young, R.A. and C.K. Mutchler. 1976. Pollution potential of manure spread on frozen ground. *J. Environ. Qual.* 5(2):174-179

Young, R.A. and R.F. Holt. 1977. Winter-applied manure: effects on annual runoff, erosion, and nutrient movement. *J. Soil Water Cons.* 32(5):219-222.



References

Zuzel, J.F., R.R. Allmaras, and R. Greenwalt. 1982. Runoff and soil erosion on frozen soils in northeastern Oregon. *J. Soil Water Cons.* 37(6):351-354.

Zuzel, J.F. and J.L. Pikul. 1987. Infiltration into a seasonally frozen agricultural soil. *J. Soil Water Cons.* 42(6):447-450.

